

TABLE 3

**Percentage distribution of women aged 18–49 at risk of unintended pregnancy, by primary contraceptive method used, according to jurisdiction, 2017\***

Jurisdiction	Female sterilization	Male sterilization	Implant	IUD	Pill	Other non-LARC hormonal	Condom	With-drawal	Other	None	Total
Alabama	19.3	4.7	†	5.4	20.6	†	17.1	†	4.2	24.1	100
Alaska	10.6	†	†	20.6	10.2	†	24.2	†	†	17.3	100
Arizona	12.3	4.1	2.2	9.3	14.7	†	16.5	1.1	6.3	32.6	100
California	9.1	1.3	3.5	9.4	19.7	3.1	22.7	2.3	5.4	23.4	100
Connecticut	7.4	3.5	2.4	8.4	21.9	1.5	20.6	†	6.1	27.2	100
Delaware	14.6	†	6.4	6.7	13.4	†	14.0	†	6.1	33.8	100
District of Columbia	3.2	†	†	9.2	13.5	†	28.9	2.9	8.6	28.9	100
Florida	12.1	3.1	5.0	7.6	15.7	1.7	20.3	0.8	6.3	27.4	100
Georgia	16.6	†	†	6.6	12.8	†	18.4	†	7.1	31.5	100
Hawaii	9.3	†	4.4	8.3	20.1	3.9	14.2	2.0	5.6	30.8	100
Idaho	10.6	4.8	†	18.1	17.7	†	15.2	†	7.3	22.5	100
Indiana	16.8	3.9	†	10.1	17.2	2.5	17.3	1.6	5.6	23.3	100
Iowa	12.8	5.1	5.2	10.4	22.7	4.7	13.8	†	3.2	21.4	100
Kansas	13.9	4.3	2.6	9.8	19.2	2.8	17.5	†	4.3	24.4	100
Louisiana	15.1	3.1	†	4.5	17.9	†	22.7	†	3.7	28.6	100
Maine	13.4	†	†	24.0	19.0	†	13.1	†	†	15.9	100
Maryland	7.9	3.4	†	11.3	18.8	2.1	23.3	†	5.6	25.6	100
Massachusetts	8.5	†	†	12.0	25.4	†	15.3	†	7.0	21.8	100
Minnesota	9.6	4.0	5.0	11.7	19.5	1.5	14.8	1.5	6.2	26.3	100
Mississippi	17.8	†	†	7.1	14.2	†	18.4	†	†	28.1	100
Missouri	19.0	4.2	2.9	7.8	17.1	†	17.2	†	5.9	24.9	100
Nebraska	12.1	4.3	3.5	8.0	18.2	†	16.5	†	5.5	28.5	100
Nevada	13.6	3.7	†	13.2	13.8	†	14.3	†	†	25.5	100
New Jersey	11.4	1.8	2.6	7.8	16.7	2.1	23.6	†	4.2	27.9	100
New Mexico	16.1	2.5	8.0	15.3	13.6	3.4	15.9	†	4.2	19.6	100
New York	8.2	1.6	†	10.2	14.7	3.1	24.6	3.1	8.1	24.4	100
North Carolina	17.2	4.3	†	9.5	17.3	3.6	15.3	†	†	22.8	100
Ohio	15.3	3.4	4.1	8.3	14.2	3.6	15.2	†	5.0	29.3	100
Oklahoma	18.7	†	†	9.0	21.5	†	12.9	†	†	26.2	100
Oregon	12.9	2.7	†	19.9	16.9	†	14.5	1.7	7.7	20.2	100
Pennsylvania	16.5	2.8	†	9.1	21.2	5.7	20.3	†	4.0	16.9	100
Puerto Rico	42.9	2.8	†	3.8	4.4	†	12.7	†	4.9	26.0	100
South Carolina	16.8	3.5	2.2	7.0	17.0	3.0	21.0	1.8	3.4	24.4	100
South Dakota	15.3	3.8	†	9.9	23.3	†	17.9	†	†	22.1	100
Texas	15.1	†	†	6.2	13.2	†	20.3	†	4.8	30.9	100
Utah	8.1	5.0	†	21.4	17.1	3.1	18.2	†	4.4	20.1	100
Virginia	12.0	2.7	4.9	10.4	18.0	2.9	20.3	†	5.8	22.1	100
West Virginia	27.4	4.9	†	8.7	14.7	†	10.1	†	4.1	25.6	100
Wisconsin	10.2	5.7	†	10.6	18.9	5.9	14.8	†	†	28.9	100
Wyoming	21.1	4.1	4.9	11.7	16.9	4.0	12.5	†	†	20.1	100

\*At risk of unintended pregnancy includes women 18–49 who were sexually active with one or more male partners, were not currently pregnant or trying to become pregnant, and had not had a hysterectomy. Women who answered “don’t care if [I] get pregnant” when asked about their rationale for nonuse were considered in this report to be at risk of unintended pregnancy (see Additional Notes on the Data for more information). †Estimate is unreliable (has a denominator of less than 50 respondents or a relative standard error greater than 30%).